

**Sequential roles for myosin-X in BMP6-dependent filopodial extension, migration, and activation of BMP receptors**

Xinchun Pi, Rongqin Ren, Russell Kelley, Chunlian Zhang, Martin Moser, Aparna B. Bohil, Melinda DiVito, Richard E. Cheney, and Cam Patterson

Vol. 179 No. 7, December 31, 2007. Pages 1569–1582.

The siRNA sequence information of myosin-X listed in the Materials and methods is incorrect. A corrected sequence is provided below. The authors apologize for this oversight.

Myo10 siRNA1 primers (5'-GATCCGCCAGACGAGAAGATATTATTCAAGAGATAAATATCTTCTCGTCTGGCTTTTTAAGCTTG-3' and 5'-AATTCAAGCTTAAAAAGCCAGACGAGAAGATATTATCTTGAATAAATATCTTCTCGTCTGGCG-3') and siRNA2 primers (5'-GATCCGCAATGCGAAGACAGTATATTCAAGAGATATACTGTCTTCGCATTGCCTTTTTAAGCTTG-3' and 5'-AATTCAAGCTTAAAAAGGC-AATGCGAAGACAGTATATCTTGAATATACTGTCTTCGCATTGCCG-3') were used for synthesis of double-stranded DNA (Invitrogen).