

## ADAPTIVE DIFFERENTIATION OF MURINE LYMPHOCYTES

### IV. (Responder $\times$ Nonresponder) F<sub>1</sub> T Cells Can Be Taught To Preferentially Help Nonresponder, Rather Than Responder, B Cells\*

By DAVID H. KATZ, LEE R. KATZ, CHERYL A. BOGOWITZ, AND PAUL H. MAURER

*From the Department of Cellular and Developmental Immunology, Scripps Clinic and Research Foundation, La Jolla, California 92037, and The Department of Biochemistry, Thomas Jefferson University, Philadelphia, Pennsylvania 19107*

Several years ago, we reported that T cells from (responder  $\times$  nonresponder) F<sub>1</sub> hybrids primed to the synthetic terpolymer, L-glutamic acid, L-lysine, L-tyrosine (GLT),<sup>1</sup> to which responses are governed by H-2-linked *Immune response-GLT* genes, were restricted in their ability to provide GLT-specific help for 2,4-dinitrophenyl (DNP)-primed B cells from the respective parental mice in response to DNP-GLT (1). Thus, such F<sub>1</sub> T cells were able to provide normal helper activity for DNP-specific B cells from responder, but not from nonresponder, donor mice. This finding contrasted sharply with the indiscriminant ability of F<sub>1</sub> T cells to interact effectively with partner B cells from either parent when the carrier antigen employed was not one to which responses were governed by a known *Ir* gene. This observation has subsequently been confirmed by others in studies conducted in mice (2, 3) and guinea pigs (4).

These observations were interpreted as an indication that in heterozygous individuals independent subpopulations of interacting T lymphocytes existed, one each corresponding to the respective parental type (1, 5, 6). Hence, we envisaged that stimulation of a (responder  $\times$  nonresponder) F<sub>1</sub> T-cell population by GLT would sensitize only the population of T cells able to recognize and react with the functional cell-interaction (CI) phenotype of the responder parent; F<sub>1</sub> T cells corresponding to the nonresponder parent CI phenotype would not be stimulated by GLT. This situation would therefore be manifested as the defective ability of F<sub>1</sub> T cells to interact with nonresponder B cells, irrespective of the antigen specificity of the latter. This original interpretation (1, 5, 6) has been reinforced by the subsequent demonstrations of the existence of independent F<sub>1</sub> T-cell subpopulations that are reactive with each respective parental CI phenotype (7-12).

\* Publication 92 from the Department of Cellular and Developmental Immunology and publication 1685 from the Immunology Departments, Scripps Clinic and Research Foundation, La Jolla, Calif. Supported by U. S. Public Health Service grants AI-13781 and AI-07825, National Foundation grants 1-540 and 1-492, American Cancer Society grant IM-5G, and Biomedical Research Support grant RRO-5514.

<sup>1</sup> Abbreviations used in this paper: ASC, *Ascaris suum* extract; CI, cell interaction; DNP, 2,4-dinitrophenyl hapten; GLT, synthetic random terpolymer of L-glutamic acid<sup>57</sup>, L-lysine<sup>38</sup>, L-tyrosine<sup>5</sup>; *Ir* gene, Immune response gene.

In this report, we present results of experiments designed to determine whether the normally restricted, cooperating phenotype of (responder  $\times$  nonresponder)  $F_1$  T cells specific for GLT could be experimentally manipulated to express a different cooperating phenotype. Specifically, we were interested in generating GLT-specific  $F_1$  T cells that might now express cooperative helper activity for nonresponder B cells. We have found that this can, indeed, be done by inducing a transient allogeneic effect during the period of priming of  $F_1$  mice to GLT. Moreover, the conditions in which the allogeneic effect is induced determines, in a critical fashion, the ultimate cooperating phenotype observed with the resulting  $F_1$  GLT-primed helper T cells.

### Materials and Methods

The terpolymer poly- $\alpha$ -(L-glutamic acid<sup>67</sup>, L-lysine<sup>38</sup>, L-tyrosine<sup>5</sup>) (GLT, lot M 72-B), mol wt 50,000, was prepared by polymerization of the  $\alpha$ -N-carboxyanhydrides of the amino acids (13). After the removal of HBr from the final lyophilized polymer, the reconstituted polymer was dissolved at pH 7.5. All other proteins, hapten-carrier conjugates, animals, methods for determining serum anti-DNP antibody levels or DNP-specific plaque-forming cells (PFC) and for the depletion of T lymphocytes by anti- $\theta$  serum plus complement, have been described (1, 14). DNP conjugates employed were DNP<sub>2,1</sub>-*Ascaris suum* (ASC) and DNP<sub>4</sub>-GLT. BALB/c, A/J, and (BALB/c  $\times$  A/J)  $F_1$ (CAF<sub>1</sub>) donors of DNP-primed B cells were primed with 10  $\mu$ g of DNP-ASC adsorbed on 4 mg of aluminum hydroxide gel (alum) and then boosted with 10  $\mu$ g of DNP-ASC in saline, 3–4 wk after primary immunization and again at monthly intervals thereafter. The mice employed in the present studies were so boosted on three occasions and finally used as B-cell donors  $\cong$  1 mo after the third saline boost. CAF<sub>1</sub> donors of GLT-primed spleen cells were immunized with 50  $\mu$ g of GLT emulsified in complete Freund's adjuvant. 10 d after this initial immunization, these mice were injected intravenously with either  $25 \times 10^6$  A/J or BALB/c spleen cells, or not injected. All mice were boosted on the same day with 50  $\mu$ g of GLT in saline; spleen cells were obtained from such donor mice 7 d after the transfer of parental cells (or none) and secondary challenge. Mixtures of primed B and T cells were transferred i.v. to 657 R irradiated CAF<sub>1</sub> recipients, which were then secondarily challenged with 50  $\mu$ g of DNP-GLT in saline i.p.

### Results

*GLT-Primed (Responder  $\times$  Nonresponder)  $F_1$  T Cells Help DNP-Primed Responder, but Not Nonresponder, Parental B Cells.* Previous studies from this laboratory, which first reported the regulatory influences of allogeneic effects on priming of carrier-specific helper T-cell activity, stressed the importance of the length of the priming regimen as well as the timing of the administration of allogeneic cells (15). Accordingly, the regimen for priming GLT-specific helper T cells in the present study was considerably abbreviated in comparison with the priming method employed in our earlier studies with this antigen (1). We felt it necessary, therefore, to investigate the patterns of cooperative helper activity manifested by CAF<sub>1</sub> helper T cells primed to GLT (in the absence of any allogeneic effects) according to the schedule used in the present studies. As shown in Fig. 1, it is quite clear that excellent cooperative secondary anti-DNP antibody responses to DNP-GLT could be obtained between  $F_1$  helper cells primed to GLT in this manner and either responder BALB/c (group IV) or  $F_1$  (group VI) B cells. In marked contrast, and consistent with our previous findings (1),  $F_1$  GLT-primed T cells were unable to provide meaningful help for nonresponder A/J B cells (group II).

*Helper T-Cell Activity of (Responder  $\times$  Nonresponder)  $F_1$  Spleen Cells Primed To GLT under the Influence of an Allogeneic Effect.* CAF<sub>1</sub> T cells obtained from mice primed to GLT, either in the absence of any allogeneic effect or in the presence of an allogeneic




FIG. 1. GLT-primed (responder × nonresponder) F<sub>1</sub> T cells help DNP-primed responder, but not nonresponder, parental B cells.

Irradiated (675 R) CAF<sub>1</sub> mice were injected i.v. with 5 × 10<sup>6</sup> anti-θ serum-treated B cells from DNP-ASC-primed A/J, BALB/c, or CAF<sub>1</sub> donors, either together with 20 × 10<sup>6</sup> GLT-primed CAF<sub>1</sub> helper T cells or without any helper cells. Helper T-cell donors were primed with 50 μg of GLT in CFA followed 10 d later by a second injection of 50 μg in saline; spleens were removed from such donor mice 7 d after the second injection to be used as helper cells. All adoptive recipients were secondarily challenged with 50 μg of DNP-GLT in saline. The data presented represent geometric means of individual numbers of IgG DNP-specific PFC per 10<sup>6</sup> recovered spleen cells of groups consisting of five mice each. Statistically significant differences in responses between experimental and control groups are indicated beside the corresponding bar.

effect induced by one or the other parental cell type, display the pattern of cooperative helper activity for DNP-primed B cells from either A/J, BALB/c, or CAF<sub>1</sub> donor mice that is depicted in Figs. 2 and 3. These data are representative of several experiments of this type, all of which utilized a standard adoptive transfer system in which varying numbers of the relevant B and T cells were transferred together to irradiated CAF<sub>1</sub> recipients. The only pertinent difference between the two experiments illustrated in these figures is that 5 × 10<sup>6</sup> B cells and 10 × 10<sup>6</sup> helper T cells were transferred in one experiment (Fig. 2), whereas 15 × 10<sup>6</sup> B cells and 30 × 10<sup>6</sup> helper T cells were transferred in the other (Fig. 3). The data is presented in terms of numbers of splenic PFC of the IgG class in Fig. 2 and levels of serum anti-DNP antibodies in Fig. 3 to illustrate the fact that the patterns of cooperative activity observed in these studies were the same, irrespective of which type of determination was used.

What these experiments have revealed is that CAF<sub>1</sub> helper T cells, primed to GLT in the absence of an allogeneic effect, provide no help for B cells from A/J mice and variable levels of help for B cells from BALB/c or CAF<sub>1</sub> donor mice depending on the numbers of GLT helper T cells transferred. Note, for example, that 10 × 10<sup>6</sup> F<sub>1</sub> helper T cells provided only marginal help for F<sub>1</sub>, and even less for BALB/c, B cells (Fig. 2), whereas significant levels of helper activity were observed in both cases when 30 × 10<sup>6</sup> GLT-primed F<sub>1</sub> cells were transferred (Fig. 3).

The remarkable findings are those pertaining to the patterns of F<sub>1</sub> helper activity, for the various B cells, that were generated under the influence of allogeneic effects




FIG. 2. Helper T-cell activity of (responder  $\times$  nonresponder) F<sub>1</sub> spleen cells primed to GLT under the influence of an allogeneic effect.

Irradiated (675 R) CAF<sub>1</sub> recipient mice were injected with i.v. with  $5 \times 10^6$  DNP-primed B cells from either A/J, BALB/c, or CAF<sub>1</sub> donors, either in the absence of helper cells or together with  $10 \times 10^6$  spleen cells taken from F<sub>1</sub> mice primed to GLT either in the same manner as described in Fig. 1, or under the influence of an allogeneic effect induced by i.v. injection (on day 10 after initial immunization with GLT) of  $25 \times 10^6$  spleen cells from parental A/J or BALB/c donors. All recipients were challenged with 50 μg of DNP-GLT in saline. The data are presented as geometric mean levels of individual IgG DNP-specific PFC/spleen of groups of five mice each assayed on day 7 after cell transfer and secondary challenge. Horizontal lines represent standard errors and relevant *P* values depicting statistically-significant differences are indicated beside the corresponding bar.

induced by one or the other parental cell type. First, note that F<sub>1</sub> helper T cells, generated under the influence of an allogeneic effect induced by either A/J or BALB/c parental cells, were clearly and significantly enhanced in their levels of cooperative activity for B cells derived from F<sub>1</sub> donors (Figs. 2 and 3, groups XI and XII). This pattern of indiscriminately enhanced F<sub>1</sub> helper T-cell activity as manifested with F<sub>1</sub> B cells did not hold true when such F<sub>1</sub> T cells were assayed for their ability to help nonresponder A/J or responder BALB/c parental B cells. Thus, F<sub>1</sub> T cells, generated under the influence of an allogeneic effect induced by parental BALB/c cells, were clearly capable of providing GLT-specific help for nonresponder A/J B cells (group IV), but were unable to provide detectable help for responder BALB/c cells (group VIII). Conversely, F<sub>1</sub> T cells, generated during an allogeneic effect induced by parental A/J cells, did not display effective helper activity for nonresponder A/J B cells (group III), although exhibiting significantly enhanced helper activity for responder BALB/c B cells (group VII).

It is pertinent to cite certain control experiments that argue against any major role of carry-over of donor parental cells into the final adoptive transfer recipients as contributing significantly to these findings. First, cell mixture experiments in which cells from F<sub>1</sub> mice primed to GLT during an allogeneic effect were analyzed for any




FIG. 3. Helper T-cell activity of (responder  $\times$  nonresponder) F<sub>1</sub> spleen cells primed to GLT under the influence of an allogeneic effect.

The same protocol as described in Fig. 2 was employed with the only difference being the numbers of transferred B and T cells given to each recipient mouse. The data are presented as geometric mean levels of serum anti-DNP antibodies of individual mice in groups of five mice each bled on day 7 after cell transfer and secondary challenge. Horizontal lines represent the range of standard errors and relevant *P* values of statistically significant differences are indicated.

possible effects on normal cooperative responses between keyhole limpet hemocyanin-primed F<sub>1</sub> T cells and parental A/J, BALB/c, or CAF<sub>1</sub> DNP-primed B cells failed to reveal any such effects. Second, analysis of such cell populations with the relevant anti-*H-2* alloantibodies has revealed little or no contamination of the F<sub>1</sub> donor cell population with detectable parental cells at the time they are removed from GLT-primed mice for the secondary adoptive transfer.

Finally, elimination of F<sub>1</sub> cells by treatment with anti-*H-2* alloantibodies plus complement directed against the opposite parental type as that used for induction of the allogeneic effect, does not leave residual cells in such F<sub>1</sub> populations capable of exerting these effects when cotransferred with mixtures of F<sub>1</sub> T and B cells such as those used in Figs. 2 and 3, group X. Representative data from such an experiment is summarized in Table I. Briefly, it can be seen that F<sub>1</sub> helper T cells, primed to GLT in the absence of an allogeneic effect, provide moderate help for DNP-primed F<sub>1</sub> B cells in adoptive recipients challenged with DNP-GLT (group II). F<sub>1</sub> helper T cells primed to GLT under the influence of an allogeneic effect, induced by parental BALB/c cells, provide enhanced levels of GLT-specific helper activities (group III). Treatment of such helper cells *in vitro* with either anti-BALB/c or anti-A/J serum + complement before transfer, abrogates the GLT-specific helper activity (groups IV and V). When equal numbers of F<sub>1</sub> GLT-primed spleen cells from donors, primed without or with the influence of an allogeneic effect, are transferred to irradiated recipients together with F<sub>1</sub> DNP-primed B cells, the magnitude of GLT-specific helper activity is essentially additive of the quantities of help exhibited by each individual

TABLE I  
*The Effects of F<sub>1</sub> Helper T Cells Primed to GLT under the Influence of an Allogeneic Effect Are Not a Result of Residual Contaminating Parental Cells\**

Group	5 × 10 <sup>6</sup> anti- $\theta$ - treated DNP- primed B cells	10 × 10 <sup>6</sup> GLT-1° CAF helper T cells	10 × 10 <sup>6</sup> GLT-1° plus allogeneic ef- fect helper T cells**	anti-DNP response† (IgG PFC/ spleen)
I		None No allogeneic effect	None	1,231
II			None	8,957
III		None	BALB/c allogeneic effect (un- treated)	15,463
IV	CAF <sub>1</sub>		BALB/c allogeneic effect (anti- BALB/c serum—Rx'd)	1,856
V			BALB/c allogeneic effect (anti-A/ J serum—Rx'd)	1,420
VI			BALB/c allogeneic effect (un- treated)	26,541
VII		No allogeneic effect	BALB/c allogeneic effect (anti- BALB/c serum—Rx'd)	9,923
VIII			BALB/c allogeneic effect (anti-A/ J serum—Rx'd)	8,796

\* Irradiated (675 R) CAF<sub>1</sub> recipients were injected i.v. with 5 × 10<sup>6</sup> DNP-primed CAF<sub>1</sub> B cells either in the absence of helper T cells or together with (a) 10 × 10<sup>6</sup> spleen cells from F<sub>1</sub> mice primed with GLT in the absence of an allogeneic effect (Fig. 1) (b) 10 × 10<sup>6</sup> F<sub>1</sub> spleen cells primed to GLT under the influence of an allogeneic effect induced by parental BALB/c cells (Fig. 2) or (c) a mixture of the two types of F<sub>1</sub> GLT-primed helper T cells. All recipients were challenged with 50 μg of DNP-GLT in saline.

\*\* F<sub>1</sub> cells primed to GLT under the influence of BALB/c parental cell-induced allogeneic effects were transferred either untreated or after treatment *in vitro* with either A/J anti-BALB/c or BALB/c anti-A/J antiserum plus complement under conditions which gave 100% lysis of each respective parental target cells.

† The data are presented as geometric mean levels of individual IgG DNP-specific PFC/spleen of groups of five mice each assayed on day 7 after cell transfer and challenge.

cell population (group VI). When helper cells from donors primed to GLT under the influence of an allogeneic effect were first pretreated with anti-BALB/c or anti-A/J serum plus complement *in vitro*, the levels of helper activity obtained reflected essentially the help provided by those cells from donors primed to GLT in the absence of any allogeneic effect (groups VII and VIII). Collectively, these data clearly document the fact that the biological activities of F<sub>1</sub> cells from donors primed to GLT under the influence of an allogeneic effect reflect function of host F<sub>1</sub> cells, and not residual parental cells used for the induction of the allogeneic effect, because alloantibodies directed against either parental cell type were equally effective in abrogating such biological activities.

### Discussion

These studies make two important points: (a) First, the usually restricted phenotype of (responder × nonresponder) F<sub>1</sub> T cells which typically is permissive only for providing cooperative helper activity for B cells of responder, but not of nonresponder, type, can be changed by inducing an allogeneic effect during the priming of such F<sub>1</sub>




FIG. 4. Influence of parental cell-induced allogeneic effect on differentiation of  $F_1$  lymphocytes. See text for explanation.

mice. This is true provided the allogeneic effect is induced by cells derived from the opposite (i.e. responder) parental type. (b) Actually, the ultimate cooperating phenotype of GLT-primed  $F_1$  helper T cells is differentially and reciprocally directed toward B cells of one parental type or the other depending on which parental donor cells are used for inducing the allogeneic effect that takes place during the priming regimen. Thus,  $F_1$  T cells, primed to GLT under influence of an allogeneic effect induced by parental BALB/c cells, now provide effective help for nonresponder A/J B cells, but do not do so for responder BALB/c B cells, and vice versa. On the other hand,  $F_1$  T cells primed to GLT under the influence of an allogeneic effect, induced by either parental cell type, display significantly enhanced levels of helper activity for B cells derived from  $F_1$  donors.

The fact that the allogeneic effect induces such exquisite discriminatory helper activities when  $F_1$  T cells are assayed on parental B cells, but loses this discriminatory aspect when  $F_1$  B cells serve as the partner B cells in the assay, is perhaps the most pertinent aspect of these findings with respect to understanding the regulatory events which these data reflect. Parenthetically, the results with  $F_1$  B cells provide additional arguments against any significant contribution made by contaminating parental B cells that may be carried over in the final assay system; but this possibility has been more directly circumvented (Results).

We believe that these results illustrate the consequences of two interdependent events which are schematically illustrated in Fig. 4: First, the allogeneic effect induced by one parental cell type exerts powerful stimulatory signals which substantially augment the normal differentiation signals induced by immunization with antigen alone; this has been amply documented to occur in many previous studies (16). The consequence of such stimulatory signals is reflected in the significant enhancement of GLT-specific helper T-cell activity provided to B cells from  $F_1$ , BALB/c, and A/J donors as shown herein. Of particular note is the capacity of the allogeneic effect, induced by BALB/c cells, to draw out permissiveness of  $F_1$  cells in providing GLT-specific help to the nonresponder A/J B cells. One point that should be clarified is that this finding is by no means directly analogous to either (a) our own earlier studies demonstrating that an appropriately timed allogeneic effect would convert the normally tolerogenic effects of DNP-derivatized D-glutamic acid, D-lysine (D-GL) to an immunogenic signal for DNP-primed B cells (17, 18) or (b) that of Ordal and Grumet (19) who demonstrated the capacity of an allogeneic effect to permit nonresponder mice to make IgG (as well as IgM) antibody responses to (T,G)-A--L.

In both of the aforementioned instances, the allogeneic effect provided a necessary stimulus to target B cells, thereby replacing a normal interaction step missing in both circumstances. In our present studies, the allogeneic effect has obviously provided the necessary stimulus, in addition to antigen, to encourage the differentiation of the subset of A/J-type T cells capable of interacting with DNP-primed A/J B cells. The reasons for the failure of such cells to differentiate through the helper cell pathway in response to GLT under normal circumstances is, of course, a central question to the whole issue of what role is actually served by *Ir* genes and their products. The present studies certainly do not provide a definitive answer to this question.

It should be pointed out, however, that the fact that the allogeneic effect was effective in inducing GLT-specific helper T cells, presumably of the nonresponder A/J-type subset, must be viewed with serious consideration in the context of the recently mounting speculation that *Ir*-gene products react specifically with antigenic determinants at the level of the macrophage to orient and display the relevant determinants for recognition by T cells of corresponding specificity (20–22). It is difficult to envisage, for example, how an allogeneic effect could circumvent an absolute requirement for *Ir*-gene-controlled antigen display by macrophages. Rather, these data seem to argue against any such absolute requirement for macrophage presentation and tend to favor the possibility that the allogeneic effect permits the development of GLT-specific  $F_1$  helper cells capable of interacting with nonresponder B cells by either (a) direct stimulation of the A/J-type subset of  $F_1$  GLT-specific T cells to differentiate into effective helper cells, or (b) elimination of some type of inherent suppressive mechanism that normally blocks development of GLT-specific helper T cells capable of cooperating with nonresponder B cells.

The second of the two interdependent events, and by far the more difficult to address, pertains to the explanation for the very striking discriminatory aspects of helper activities when  $F_1$  T cells are assayed on parental B cells. Obviously, no direct evidence is presently available to permit conclusions concerning this aspect of our data. Nevertheless, we would like to suggest that aside from the stimulatory consequences of the allogeneic effect in inducing helper T-cell function as discussed above, a second consequence may be the stimulation of an  $F_1$  response against certain of its own receptors which self-recognize native CI determinants ( $\alpha$ -CI). Thus, as illustrated in Fig. 4, when parental B cells induce an allogeneic effect in an ( $A \times B$ )  $F_1$ , the response within the  $F_1$  would be directed against self receptors for B-type CI molecules (anti- $\alpha$  CI<sub>B</sub>), and vice-versa. These types of anti-idiotypic responses against self-recognizing receptors would be capable of preventing development of helper T cells belonging to the corresponding parental-type T cell subpopulation without adversely affecting the development of helper cells corresponding to the opposite parental-type T-cell subpopulation.

In a sense, these postulated anti- $\alpha$  CI responses are analogous to, but not necessarily the same as, the  $\alpha$ -major histocompatibility complex (MHC) idiotypic reactions described recently by Bellgrau and Wilson (23) which confer specific resistance to parental cell-induced graft-vs.-host disease in irradiated  $F_1$  rats.

If studies currently in progress demonstrate anti- $\alpha$  CI reactions to be the mechanism responsible for the data presented here, then one wonders (a) if responses of this type occur normally in the pathways of immune regulation, (b) whether virgin cells and primed cells are similarly susceptible to the effects of such reactions, or not, and (c)

what relationship such anti- $\alpha$  CI responses might have to the whole picture portrayed by *Ir* genes and the specificity they appear to display for the antigens under their control.

### Summary

Responses to the synthetic terpolymer L-glutamic acid, L-lysine, L-tyrosine (GLT) in the mouse are controlled by *H-2*-linked *Ir-GLT* genes. (Responder  $\times$  nonresponder)  $F_1$  hybrid mice, themselves phenotypic responders, can be primed with GLT to develop specific helper cells capable of interacting with 2,4-dinitrophenyl hapten (DNP)-primed  $F_1$  B cells in response to DNP-GLT. Unlike the indiscriminant ability of  $F_1$  helper T cells for conventional antigens (i.e. not *Ir* gene-controlled), which can help B cells of either parental type (as well as  $F_1$ ) equally well, GLT-primed  $F_1$  T cells can only provide help under normal circumstances for B lymphocytes of responder parent origin; they are unable to communicate effectively with nonresponder parental B cells (1, and the present studies). The present studies reveal, however, that the induction of a parental cell-induced allogeneic effect during priming of  $F_1$  mice to GLT actually dictates the direction of cooperating preference that will be displayed by such  $F_1$  helper cells for B cells of one parental type or the other. Thus,  $F_1$  T cells, primed to GLT under the influence of an allogeneic effect induced by parental BALB/c cells, develop into effective helpers for nonresponder A/J B cells, but fail to develop effective helpers for responder BALB/c B cells, and vice-versa. In contrast,  $F_1$  T cells, primed to GLT under the influence of an allogeneic effect induced by either parental type, display significantly enhanced levels of helper activity for B cells derived from  $F_1$  donors. These results are interpreted to reflect the existence of two interdependent events provoked by the allogeneic effect: one event augments the differentiation of GLT-specific helper T cells belonging to the subset corresponding to the opposite parental type; this would explain the development of increased helper activity provided to partner B cells of opposite parental type (as well as of  $F_1$  origin). The second event, we postulate, involves the production of responses against the receptors which normally self-recognize native cell interaction determinants; this form of anti-idiotypic response is restricted against self-recognizing receptors of the same parental type used for induction of the allogeneic effect, hence explaining diminished helper activity of such  $F_1$  cells for partner B lymphocytes of corresponding parental type.

We are very grateful to Norman Klinman and Hiroshi Yamamoto for critically reviewing the manuscript and offering helpful suggestions. The capable assistance of Anthea Hugus and Keith Dunn in the preparation of this manuscript is greatly appreciated.

*Received for publication 27 December 1979.*

### References

1. Katz, D. H., T. Hamaoka, M. E. Dorf, P. H. Maurer, and B. Benacerraf. 1973. Cell interactions between histoincompatible T and B lymphocytes. IV. Involvement of the immune response (*Ir*) gene in the control of lymphocyte interactions in response controlled by the gene. *J. Exp. Med.* **183**:734.
2. Pierce, S. K. 1977. Recognition restrictions in lymphocyte collaborative interactions in  $IgG_1$

- antibody responses. *In Immune System: Genetics and Regulation*. E. E. Sercarz, L. A. Herzenberg, and C. F. Fox, editors. Academic Press, Inc., New York. 447.
3. Press, J. L., and H. O. McDevitt. 1977. Allotype-specific analysis of anti-(Tyr,Glu)-Ala-Lys antibodies produced by Ir-IA high and low responder chimeric mice. *J. Exp. Med.* **146**:1815.
  4. Yamashita, U., and E. M. Shevach. 1978. The histocompatibility restrictions on macrophage T-helper cell interaction determine the histocompatibility restrictions on T-helper cell B-cell interaction. *J. Exp. Med.* **148**:1171.
  5. Katz, D. H., and B. Benacerraf. 1974. The role of histocompatibility gene products in cooperative cell interactions between T and B lymphocytes. *In The Immune System: Genes, Receptors, Signals*. E. E. Sercarz, A. R. Williamson, and C. Fred Fox, editors. Academic Press, Inc., New York. 569.
  6. Katz, D. H. 1977. The role of the histocompatibility gene complex in lymphocyte differentiation. *Cold Spring Harbor Symp. Quant. Biol.* **41**:611.
  7. Skidmore, B. J., and D. H. Katz. 1977. Haplotype preference in lymphocyte differentiation. I. Development of haplotype-specific and suppressor activities in F<sub>1</sub> hybrid activated T cell populations. *J. Immunol.* **119**:694.
  8. Paul, W. E., E. M. Shevach, D. W. Thomas, S. F. Pickeral, and A. S. Rosenthal. 1977. Genetic restriction in T-lymphocyte activation by antigen-pulsed peritoneal exudate cells. *Cold Spring Harbor Symp. Quant. Biol.* **41**:571.
  9. Miller, J. F. A. P., and M. A. Vadas. 1977. The major histocompatibility complex: Influence on immune reactivity and T-lymphocyte activation. *Scand. J. Immunol.* **6**:771.
  10. Thomas, D. W., and E. M. Shevach. 1978. Nature of the antigenic complex recognized by T lymphocytes. V. Genetic predisposition of independent F<sub>1</sub> T cell subpopulations responsive to antigen-pulsed parental macrophages. *J. Immunol.* **120**:638.
  11. McDougal, J. S., and S. P. Cort. 1978. Generation of T helper cells *in vitro*. IV. F<sub>1</sub> T helper cells primed with antigen-pulsed parental macrophages are genetically restricted in their antigen-specific helper activity. *J. Immunol.* **120**:445.
  12. Swierkosz, J. E., K. Rock, P. Marrack, and J. W. Kappler. 1978. The role of *H-2*-linked genes in helper T-cell function. II. Isolation on antigen-pulsed macrophages of two separate populations of F<sub>1</sub> helper T cells each specific for antigen and one set of parental *H-2* products. *J. Exp. Med.* **147**:554.
  13. Katchalski, E., and M. Sela. 1958. Synthesis and chemical properties of poly- $\alpha$ -amino acids. *Adv. Protein Chem.* **13**:243.
  14. Katz, D. H., and D. P. Osborne, Jr. 1972. The allogeneic effect in inbred mice. II. Establishment of the cellular interactions required for enhancement of antibody production by the graft-versus-host reaction. *J. Exp. Med.* **136**:455.
  15. Osborne, D. P., Jr., and D. H. Katz. 1973. The allogeneic effect in inbred mice. IV. Regulatory influences of graft-versus-host reactions on host T lymphocyte functions. *J. Exp. Med.* **138**:825.
  16. Katz, D. H. 1972. The allogeneic effect on immune responses. Model for regulatory influences of T lymphocytes on the immune system. *Transplant. Rev.* **12**:141.
  17. Katz, D. H., J. M. Davie, W. E. Paul, and B. Benacerraf. 1971. Carrier function in anti-hapten antibody responses. IV. Experimental conditions for the induction of hapten-specific tolerance or for the stimulation of anti-hapten anamnestic responses by "non-immunogenic" hapten-polypeptide conjugates. *J. Exp. Med.* **134**:201.
  18. Osborne, D. P., Jr., and D. H. Katz. 1973. The allogeneic effect in inbred mice. III. Unique antigenic structural requirements in the expression of the phenomenon on unprimed cell populations *in vivo*. *J. Exp. Med.* **137**:991.
  19. Ordal, J. C., and F. C. Grumet. 1972. Genetic control of the immune response. The effect of graft-versus-host reaction on the antibody responses to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys in nonresponder mice. *J. Exp. Med.* **136**:1195.

20. Rosenthal, A. S. 1978. Determinant selection and macrophage function in genetic control of the immune response. *Immunol. Rev.* **40**:136.
21. Benacerraf, B. 1978. A hypothesis to relate the specificity of T lymphocytes and the activity of I region-specific Ir genes in macrophages and B lymphocytes. *J. Immunol.* **120**:1809.
22. Schwartz, R. H. 1978. A clonal deletion model for Ir gene control of the immune response. *Scand. J. Immunol.* **7**:3.
23. Bellgrau, D., and D. B. Wilson. 1978. Immunological studies of T cell receptors. I. Specifically induced resistance to graft-versus-host disease in rats mediated by host T-cell immunity to alloreactive parental T cells. *J. Exp. Med.* **148**:103.