

HUMAN RHEUMATOID FACTOR CROSSIDIOTYPES
II. Primary Structure-dependent Crossreactive Idiotypic, PSL2-CRI,
Present on Wa Monoclonal Rheumatoid Factors Is Present on Bla and
Other IgM κ Monoclonal Autoantibodies

BY VINCENT AGNELLO,* FERNANDO GOÑI,‡ J. L. BARNES,*
MARIA-TERESA DE LA VEGA,‡ AND BLAS FRANGIONE‡

*From the *Department of Laboratory Medicine, Lahey Clinic Medical Center,
Burlington, Massachusetts 01805; and the Edith Nourse Rogers Memorial Veterans
Administration Hospital, Bedford, Massachusetts 01730; and the ‡Department of Pathology,
Kaplan Cancer Center, New York University Medical Center, New York 10016*

The major human monoclonal rheumatoid factor (mRF) crossidiotypic (XId) Wa is present on ~60% of mRFs (1). Distinct from Wa are the two minor groups, Po and Bla (1, 2). The three XIDs were initially defined serologically using polyclonal antisera raised to native mRFs. Subsequently, by using antisera raised to synthetic hypervariable region (complementarity determining region [CDR]) peptides of a Wa group mRF (Sie), a primary structure-dependent XId was defined that was thought to be unique to the Wa group (3–5). Analysis of light chain amino acid sequences of Wa mRFs revealed that only the second CDR (L-CDR2) was identical in all groups. The crossreactive idiotype (PSL2-CRI) was defined using an antiserum to the second CDR peptide of the Sie light chain (anti-PSL2). Anti-PSL2 reacted with all Wa mRFs tested and only weakly and not at all, respectively, with the proteins Pom and Lay of the Po group. These and other data obtained were consistent with the possibility that the PSL2-CRI was unique to the Wa mRFs. However, in a study of amino acid sequences of κ IIIb light chains of monoclonal human IgM κ autoantibodies, an antibody with specificity for low-density lipoprotein (Son) had the same L-CDR2 amino acid sequence as the Wa mRFs (6). This protein did not have anti- γ -globulin activity and did not bear the Wa XId (7). These data suggested that the PSL2-CRI may not be confined to Wa mRFs. The third mRF XId group (Bla) was not investigated in the initial PSL2-CRI studies. Recent study (8) has demonstrated the presence of PSL2-CRI on Bla mRF, the prototype of the Bla group. The present study was undertaken to determine the amino acid sequence of the L-CDR2 of Bla, to compare it with the L-CDR2 sequence of Wa and Po mRFs, and to determine whether the PSL2-CRI is present on other monoclonal autoantibodies. Preliminary reports (9, 10) of these studies have been published.

This work was supported by National Institutes of Health grants AM-35487 (to V. Agnello) and AM-02594 (to B. Frangione), and Veterans Administration Merit Review Grant. Fernando Goñi is a Fellow of the Arthritis Foundation, Atlanta, GA. Address correspondence to Vincent Agnello, Department of Laboratory Medicine, Lahey Clinic Medical Center, 41 Mall Road, Burlington, MA 01805.

Materials and Methods

The isolation and idiotypic characterization of the IgM κ mRFs Wa and McD of the Wa group, Lay of the Po group, and Bla of the Bla group have been described in detail previously (1, 2). Isolation of the IgM κ monoclonal anti-low-density lipoprotein (Son) and analysis of light chain amino acid sequences have also been reported previously (6). Sera of Joh and Gag, IgM κ monoclonal cold agglutinins, were provided by Dr. R. Williams (Univ. of New Mexico Hospital, Albuquerque, NM). Partial analyses of the amino acid sequences of the light chains of these two proteins have been performed previously, and both characterized as V κ III (11). The monoclonal cold agglutinins were isolated from serum by high-pressure liquid chromatography using 5000-PW column 7.5 mm \times 30 cm (Altex Spherogel-TSK; Beckman Instruments, Inc., Fullerton, CA). Immunoblots of denatured light chains isolated by SDS-PAGE were tested by probing Western blots of the gels as reported (8). Anti-PSL2 and anti-PSL3 (4) were provided by Drs. P. P. Chen and D. A. Carson (Scripps Clinic, La Jolla, CA). Goat anti-human κ chain serum was obtained from Kallestad Laboratories (Austin, TX). Enzymatic digestion, purification of peptides, and analysis of amino acid sequences were performed as described (12).

Results and Discussion

The amino acid sequence of the L-CDR2 of the Bla protein shown in Fig. 1 was identical to that reported for the Wa mRFs. The reactivity of anti-PSL2, anti-PSL3, and anti- κ with denatured light chain of the various monoclonal autoantibodies is shown in Fig. 2. Anti-PSL2 reacted with all light chains except those from Lay, the Po mRF. Anti-PSL3 reacted with light chains of the Wa mRFs, Wa and McD, and Son, the anti-low-density lipoprotein, but not with the light chains of Bla or the cold agglutinins.

The demonstration in this study that the L-CDR2 of Bla mRF is identical to that of the Wa mRFs and that the PSL2-CRI, as determined by anti-PSL2, is present on Bla, a mRF that reacts with DNA nucleoprotein, Son, a monoclonal anti-low-density lipoprotein, and two monoclonal cold agglutinins unequivocally rules out that the PSL2-CRI is associated only with mRFs of the Wa group. On the contrary, these findings suggest that the PSL2-CRI may be a common determinant on most IgM κ monoclonal autoantibodies and raise the possibility that the PSL2-CRI is an idiotype involved in a more general regulation of autoantibodies. The high incidence of PSL2-CRI among rheumatoid factors in essential mixed cryoglobulinemia, a disorder in which the rheumatoid factors are usually monoclonal, also supports the hypothesis that PSL2-CRI is associated with monoclonal IgM κ autoantibodies (4).

A major alternative possibility to that of selective association of PSL2-CRI with monoclonal autoantibodies is that the presence of PSL2-CRI among monoclonal autoantibodies reflects the occurrence of the PSL2-CRI in the IgM κ pool in pathologic sera. This must be considered for several reasons. A high incidence of PSL2-CRI occurs among V κ IIIb light chains (4, 6); V κ IIIb light chains make up a substantial portion of κ light chains of polyclonal IgM in normal serum, whereas only trace amounts are present among light chains of IgG and IgA (13). Uncertainty exists about the percentage of V κ IIIb light chains present among IgM κ proteins (13, 14) and about the effect of polyclonal increase of IgM on V κ IIIb concentrations (15). Therefore, the high incidence of PSL2-CRI among monoclonal autoantibodies may reflect the relatively high levels of IgM κ IIIb that may be present in pathologic sera.

	CDR2						
	Amino Acid Position						
	50	51	52	53	54	55	56
Bla	Gly	Ala	Ser	Ser	Arg	Ala	Thr
Wa group	_____						
Wof	_____						
Sie	_____						
Gar	_____						
Got	_____						
Neu	_____						
Flo	_____						
Po group	_____						
Lay				Thr		Glu	Ala

FIGURE 1. Bla light chain CDR2 amino acid sequence compared with reported Wa and Po mRF CDR2 sequences (12).




FIGURE 2. Triplicate immunoblots of denatured light chains from Bla mRF, Lay mRF, Wa mRF, Son anti-low-density lipoprotein monoclonal antibody, Joh monoclonal cold agglutinin, Gag monoclonal cold agglutinin, and McD mRF. Blots are probed with anti-PSL2, anti-PSL3, or anti- κ , then with ^{125}I -labeled protein A.

Whether $V\kappa\text{IIIb}$ and the PSL2-CRI are selectively present in autoantibodies could possibly be determined more definitively from studies of polyclonal autoimmune disease states. Thus far, studies on polyclonal rheumatoid factors are conflicting. Qualitative studies have shown a high incidence of PSL2-CRI, whereas quantitative studies for $V\kappa\text{IIIb}$ indicate that only small amounts are present (16, 17). From the evidence presented here and previous studies, which demonstrated that the Wa and Bla XIDs are conformational antigens requiring both heavy and light chains, it is clear that the determinant detected by anti-PSL2 is primary structure antigen not directly related to the Wa or Bla XID determinants. It remains to be determined whether PSL2-CRI is indeed an XID that is selectively present on autoantibodies or an allotypic marker for a $V\kappa\text{III}$ germline gene. The latter has not been excluded because somatic mutation of the L-CDR2 could explain the presence of $V\kappa\text{III}$ light chains that do not react with anti-PSL2.

Summary

The amino acid sequence of the L-CDR2 (complementarity-determining region) of Bla mRF (monoclonal rheumatoid factor) is identical to that of the Wa mRFs. The PSL2-CRI (crossreactive idiotype), as determined by anti-PSL2, which has been shown to be present on all Wa mRFs, is also present on the Bla mRF and other monoclonal autoantibodies. PSL2-CRI is, therefore, not unique to Wa mRFs and may be present on most IgM κ monoclonal autoantibodies. Whether PSL2-CRI is a crossidiotype (XId) that is selectively present on autoantibodies or represents an allotypic marker for a V κ III gene is undetermined.

We thank Ms. Patricia Leasure for excellent technical assistance, Drs. Pojen Chen and Dennis Carson for kindly providing anti-PSL2 and anti-PSL3, and Dr. Ralph Williams for kindly providing the monoclonal cold agglutinins.

Received for publication 15 September 1986.

References

1. Kunkel, H. G., V. Agnello, F. G. Joslin, R. J. Winchester, and J. D. Capra. 1973. Cross-idiotypic specificity among monoclonal IgM proteins with anti- γ -globulin activity. *J. Exp. Med.* 137:331.
2. Agnello, V., A. Arbetter, G. I. de Kasep, R. Powell, E. M. Tan, and F. Joslin. 1980. Evidence for a subset of rheumatoid factors that cross-react with DNA-histone and have a distinct cross-idiotype. *J. Exp. Med.* 151:1514.
3. Chen, P. P., S. Fong, D. Normansell, R. A. Houghten, J. G. Karras, J. H. Vaughan, and D. A. Carson. 1984. Delineation of a crossreactive idiotype on human autoantibodies with antibody against a synthetic peptide. *J. Exp. Med.* 159:1502.
4. Chen, P. P., F. Goñi, R. A. Houghten, S. Fong, R. Goldfien, J. H. Vaughan, B. Frangione, and D. A. Carson. 1985. Characterization of human rheumatoid factors with seven antiidiotypes induced by synthetic hypervariable region peptides. *J. Exp. Med.* 162:487.
5. Chen, P. P., F. Goñi, S. Fong, F. Jirik, J. H. Vaughan, B. Frangione, and D. A. Carson. 1985. The majority of human monoclonal IgM rheumatoid factors express a "primary structure-dependent" cross-reactive idiotype. *J. Immunol.* 134:3281.
6. Pons-Estel, B., F. Goñi, A. Solomon, and B. Frangione. 1984. Sequence similarities among κ IIIb chains of monoclonal human IgM κ autoantibodies. *J. Exp. Med.* 160:893.
7. Posnett, D. N., R. Wisniewolski, B. Pernis, and H. G. Kunkel. 1986. Dissection of human antigammaglobulin idiotype system with monoclonal antibodies. *Scand. J. Immunol.* 23:169.
8. Agnello, V., and J. L. Barnes. 1986. Human rheumatoid factor cross-idiotypes: I. WA and BLA are heat-labile conformational antigens requiring both heavy and light chains. *J. Exp. Med.* 164:1809.
9. Agnello, V., J. L. Barnes, F. Goñi, and B. Frangione. 1986. The primary structure cross idiotype PSL2-XId associated with WA monoclonal rheumatoid factors is present on other monoclonal autoantibodies. *Proc. IXth Pan Am. Congr. Rheumatol. Buenos Aires*, 9.
10. Agnello, V., F. Goñi, J. L. Barnes, and B. Frangione. 1986. Cross-idiotypes of human monoclonal rheumatoid factors. *Proc. 6th Internat. Congr. Immunol.* 3:73.10.
11. Capra, J. D., J. M. Kehoe, R. C. Williams, Jr., T. Feizi, and H. G. Kunkel. 1972. Light chain sequences of human IgM cold agglutinins. *Proc. Natl. Acad. Sci. USA.* 69:40.

12. Goñi, F., P. P. Chen, B. Pons-Estel, D. A. Carson, and B. Frangione. 1985. Sequence similarities and cross-idiotypic specificity of L chains among human monoclonal IgM κ with anti- γ -globulin activity. *J. Immunol.* 135:4073.
13. Greenstein, J. L., and G. N. Abraham. 1984. V κ IIIb expression is altered in acquired and drug-induced hypogammaglobulinemia. *Immunology.* 53:779.
14. Moynihan, J. A., R. J. Looney, and G. N. Abraham. 1985. The V κ IIIb light chain subgroup: restricted association with mu heavy chain in normal human serum. *Immunology.* 54:207.
15. Moynihan, J. A., R. J. Looney, and G. N. Abraham. 1986. The V κ IIIb light chain sub-subgroup. II. Isotype expression in acquired hypogammaglobulinaemia and as a function of age. *Immunology.* 57:325.
16. Fong, S., P. P. Chen, T. A. Gilbertson, J. R. Weber, R. I. Fox, and D. A. Carson. 1986. Expression of three cross-reactive idiotypes on rheumatoid factor autoantibodies from patients with autoimmune diseases and seropositive adults. *J. Immunol.* 137:122.
17. Williams, J. M., P. D. Gorecic, R. J. Looney, and G. N. Abraham. 1986. Association of V κ IIIb immunoglobulin light chains with IgM anti-IgG autoantibodies. *Arthritis Rheum.* 29:S33.