

Preferential Usage of the Fc Receptor γ Chain in the T Cell Antigen Receptor Complex by γ/δ T Cells Localized in Epithelia

By Hiroshi Ohno,* Satoru Ono,* Nakami Hirayama,†
Shinji Shimada,§ and Takashi Saito*

From the *Division of Molecular Genetics, Center for Biomedical Science, Chiba University School of Medicine, Chiba 260; the †Department of Immunology, National Institute of Health, Tokyo 162; and the §Department of Dermatology, Tokyo University Branch Hospital, Tokyo 112, Japan

Summary

ζ and η chains of the T cell antigen receptor (TCR) complex and the γ chain of Fc receptors (FcR γ) constitute a family of proteins important for the expression of, and signal transduction through, these receptors in hematopoietic cells. In ζ -deficient mice, TCR expression was reduced in most T cells. By contrast, CD8 $\alpha\alpha^+$ TCR- γ/δ^+ intestinal intraepithelial lymphocytes in these mice expressed a normal level of TCR. Biochemical analysis of the TCR complex in these cells from ζ -deficient as well as normal mice revealed the predominant usage of FcR γ . Furthermore, γ/δ^+ T cells in epithelia of the skin and female reproductive organs from ζ -deficient mice also showed relatively high TCR expression, indicating the usage of FcR γ . These observations demonstrate the preferential usage of FcR γ by γ/δ^+ T cells localized in epithelia of normal mice.

T cells recognize antigen bound to the products of MHC on APC through TCR, and are activated to exert various effector functions (1, 2). The TCR complex is a multimeric complex composed of three groups of proteins; the TCR- α/β (or $-\gamma/\delta$) dimer, the CD3 complex, and the ζ family of disulfide-linked dimers (3). TCR dimers are responsible for antigen binding, whereas the other molecules are thought to be important for transmembrane signaling.

The ζ family has three known members, ζ , η , and FcR γ (4–6). In thymocytes and peripheral T cells, most of TCR complexes contain ζ homodimers, whereas 5–10% of TCRs associate with the ζ - η heterodimer. On the other hand, the TCR complex in a minor population of T cells has ζ and η as a form of heterodimers with FcR γ (4, 5). Moreover, it is reported that FcR γ was exclusively associated with TCR in in vitro cultured large granular lymphocytes (6).

A considerable body of evidence shows the importance of ζ both in assembly and surface expression of the TCR complex and in receptor-mediated signal transduction (7–13). Cytoplasmic tails of ζ family as well as CD3 molecules have consensus motifs important for intracellular signal transduction (14, 15). Recent studies indicate that the TCR complex has two signal transduction modules, the ζ family dimer and the CD3 complex, and that the two modules could transduce distinct signals into the cells (15, 16). In addition, it has been reported that the TCR complex containing FcR γ has a different signaling capacity from those containing ζ

(17–19). Consequently, differential usage of ζ family molecules by distinct subsets of T cells may reflect difference in lineage and/or function of these cells.

Recently, we and others demonstrated that ζ is critical for normal T cell development and function, using mice deficient in the expression of ζ (20, 21). We also showed that η is not so efficient in assembly and surface expression of TCR complexes as ζ (20). Biochemical analysis of T hybridoma cells revealed that this was caused by retention in the endoplasmic reticulum of η -containing TCR complexes (22). Taking advantage of the mice lacking ζ (ζ^0 mice) (20), we evaluated FcR γ usage by various subsets of T cells.

Materials and Methods

Mice. C57BL/6 and KSN *nu/nu* mice (23) were purchased from Japan SLC, Inc. (Hamamatsu, Japan). ζ^0 mice (20) were bred in our facility.

Cell Preparation. Single cell suspensions of thymocytes and splenocytes were prepared in RPMI 1640 supplemented with 10% heat-inactivated FCS, 2 mM glutamine, 50 μ M 2-ME, and 100 μ g/ml kanamycin. Splenocytes were depleted of erythrocytes by lysis. For enrichment of CD4⁻8⁻ double-negative (DN) thymocytes, thymocytes were incubated with culture supernatant of anti-CD4 mAb (MT4), plated on plastic dishes precoated with rabbit anti-mouse immunoglobulin (Cappel Laboratories, Cochranville, PA), and nonadherent cells were collected as DN thymocytes-enriched population. Intestinal intraepithelial lymphocytes (i-IEL), hepatic lymphocytes, lymphocytes in mucosal epithelia of




Figure 1. Expression of TCR- α/β and - γ/δ dimers on thymocytes, splenocytes, and i-IEL from $\zeta T/\zeta T$ mice. (A) CD3 ϵ and TCR β or TCR δ expression on the surface of thymocytes, splenocytes, and i-IEL were examined as described in Materials and Methods. (B) CD4, CD8, and TCR δ expression on the surface of i-IEL were analyzed. CD4/CD8 α or CD8 α /CD8 β were used to divide i-IEL into four subpopulations (*top*). The histograms shown below represent the fluorescence intensity for TCR- δ detected by the third fluorescence, within each quadrant.

the female reproductive tract (r-IEL), and Thy-1⁺ dendritic epidermal T cells (DETC) were isolated as previously described (24–27).

Flow Cytometric Analysis. Cells from the thymus, spleen, peripheral blood, liver, i-IEL, DETC, and r-IEL were stained with mAbs recognizing CD3 ϵ , TCR- α/β , TCR- γ/δ , CD4, CD8 α , CD8 β , and IL-2R β , and the surface expression of these molecules was analyzed on a FACScan[®] flow cytometer (Becton Dickinson & Co., Mountain View, CA). The following mAbs were used: PE-conjugated anti-CD4 (GK1.5) and FITC-labeled anti-CD8 α (53-6.7), purchased from Becton Dickinson & Co., FITC-conjugated or biotinylated anti-CD3 ϵ (145-2C11); biotinylated anti-TCR- β (H57-597), biotinylated anti-TCR- γ/δ (GL3); and PE-labeled anti-CD45R/B220 (RA3-6B2) from PharMingen (San Diego, CA); biotinylated anti-IL-2R β (TM- β 1) was kindly provided by Dr. M. Miyasaka (Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan). Biotinylated mAbs were developed with streptavidin-PE (Becton Dickinson & Co.) or streptavidin-Tri-Color[®] (Caltag Laboratories, San Francisco, CA). Dead cells were excluded by staining with propidium iodide. Cells in the lymphocyte gate defined by light scatter were collected.

Cell Surface Biotinylation, Immunoprecipitation, and Two-dimensional SDS-PAGE Analysis. Cell surface biotinylation was performed as

previously described (28). Cells were then solubilized in lysis buffer (1% digitonin, 50 mM Tris-HCl, pH 7.6, 300 mM NaCl, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin, 1 mM PMSF, 10 mM iodoacetamide) at a concentration of 10⁸ cells/ml. Immunoprecipitation was performed with anti-TCR- δ mAb (3A10), anti-TCR- β mAb (H57-597), anti- ζ mAb (H146.698-A), and anti-Fc γ antiserum, which were kindly provided by Drs. S. Tonegawa (Massachusetts Institute of Technology, Boston, MA), R. Kubo (Cytel Corporation, San Diego, CA), and C. Ra (Juntendo University, Tokyo, Japan), respectively. Immunoprecipitates were resolved by two-dimensional nonreducing-reducing SDS-PAGE (14% for the first dimension and 16% for the second dimension), transferred onto polyvinylidene fluoride membrane (Immobirom-P[®]; Millipore Corp., Bedford, MA). Membranes were soaked in skim-milk in PBS, and biotinylated proteins were detected using streptavidin-peroxidase (VECTA-STAIN[®] Elite ABC kit; Vector Laboratories Inc., Burlingame, CA) and ECL[®] system (Amersham International, Buckinghamshire, England).

Results and Discussion

The expression levels of TCR on thymocytes and peripheral T cells were significantly reduced in $\zeta T/\zeta T$ mice (20).




Figure 2. Biochemical analysis of the surface TCR composition. (A) Total and (B) DN-enriched thymocytes from wild type mice, i-IEL from (C and D) C57BL/6 (+/+) mice, (E) KSN *nu/nu* mice, and (F) ζ T/ ζ T mice were surface biotinylated, lysed, and the TCR complex was immunoprecipitated with (A and C) anti-TCR- β mAb (H57-597), or B, D, E, and F) anti-TCR- δ mAb (3A10). Immunoprecipitates were separated by two-dimensional nonreducing-reducing SDS-PAGE, transferred onto PVDF membrane, and biotinylated proteins were detected as described in Materials and Methods. Positions of ζ - ζ (\uparrow), ζ -FcR γ (\blacktriangle), and FcR γ -FcR γ (\triangle) dimers are indicated. Note that heterodimers of p12 (35, and Kuwabara, I., and Saito, T., manuscript in preparation) with ζ and FcR γ are observed in some panels. In three experiments (including E), densitometric analysis showed that the mean of the FcR γ of ζ -FcR γ dimers/FcR γ of FcR γ -FcR γ dimers is 1:11.9. Considering that the molar ratio of ζ and FcR γ in ζ -FcR γ dimers is 1, the expression level of FcR γ molecules is 13 times more than that of ζ .

This was observed in both α/β^+ and γ/δ^+ thymocytes as well as splenic T cells (Fig. 1 A). These data indicate that those cells predominantly express ζ as a component of the TCR complex, and extended to normal T cells the previous observation on in vitro cell lines that the expression of ζ is critical for a normal level of surface TCR expression (7-10).

When we analyzed i-IEL, however, a striking difference was observed (Fig. 1 A) (20). Similar to thymocytes and peripheral T cells, surface TCR expression of α/β^+ i-IEL was impaired by the ζ T mutation. By contrast, surface TCR level of γ/δ^+ i-IEL from ζ T/ ζ T mice was comparable with that of γ/δ^+ i-IEL from wild type mice. In γ/δ^+ i-IEL, two distinct subpopulations are known; the major population of CD8 $\alpha\alpha$ -bearing cells and the minor DN cells. The former is thought to differentiate extrathymically, whereas the latter is believed to derive from the thymus (29). Whether TCR expression of both γ/δ^+ i-IEL populations was resistant to the ζ T mutation was analyzed by three-color flow cytometry. As shown in Fig. 1 B, the expression level of surface TCR in CD8 $\alpha\alpha^+$ γ/δ^+ i-IEL from ζ T/ ζ T mice remained normal, but that in DN γ/δ^+ i-IEL was reduced. These observations suggest that, in CD8 $\alpha\alpha^+$ γ/δ^+ i-IEL, ζ family molecules other than ζ were predominantly associated with TCR.

To test this possibility, various preparations of T cells were analyzed for their composition of TCR complexes by a sensitive surface biotinylation (28) and two-dimensional non-reducing-reducing SDS-PAGE. TCR on α/β^+ and γ/δ^+ thymocytes as well as α/β^+ i-IEL from wild type mice was

predominantly associated with ζ - ζ homodimers (Fig. 2, A-C). By contrast, analysis of TCR complexes on γ/δ^+ i-IEL revealed that, on these cells, both ζ and FcR γ contribute for TCR constitution (Fig. 2 D). Identification of ζ and FcR γ was confirmed by direct precipitation of these molecules with anti- ζ mAb (H146.968A) and anti-FcR γ antiserum, respectively (data not shown). Existence of two subpopulations in γ/δ^+ i-IEL from normal mice as mentioned above may complicate the analysis. To avoid this complexity, we utilized γ/δ^+ i-IEL from nude mice, which consist exclusively of CD8 $\alpha\alpha^+$ cells (Ohno, H., unpublished observation). The result was more striking; these cells predominantly utilize FcR γ as TCR subunits, and ζ homodimers were hardly seen (Fig. 2 E). Assuming that the labeling efficiency of FcR γ is comparable between FcR γ homodimers and FcR γ - ζ heterodimers, these cells express 13 times more FcR γ molecules than ζ in TCR (see legend to Fig. 2). It is likely that DN γ/δ^+ i-IEL, like γ/δ^+ thymocytes, predominantly utilize ζ based on the following reasons: the amount of precipitated ζ from γ/δ^+ i-IEL from normal mice was much higher than that from nude mice; and TCR expression in DN γ/δ^+ i-IEL from ζ T/ ζ T mice was reduced. We further examined whether the predominance of FcR γ in CD8 $\alpha\alpha^+$ γ/δ^+ i-IEL is also observed in ζ T/ ζ T mice. Fig. 2 F shows that this was the case. Collectively, these data indicate that the normal level of TCR expression in CD8 $\alpha\alpha^+$ γ/δ^+ i-IEL from ζ T/ ζ T mice correlates the predominant usage of FcR γ by TCR in this subset.

The preferential usage of FcR γ as a TCR component in




Figure 3. Expression of TCR on hepatic lymphocytes, r-IEL, and DETC from $\zeta T/\zeta T$ mice. (A) CD3 ϵ and IL-2R β expression on hepatic lymphocytes was analyzed as described in Materials and Methods. CD3^{int}IL-2R β ^{hi} cells, which are thought to develop extrathymically in the liver (30), are indicated with squares. (B) CD3 ϵ and TCR δ expression on peripheral blood cells, r-IEL, and DETC were examined as described in Materials and Methods.

CD8 $\alpha\alpha^+$ TCR γ/δ^+ i-IEL propelled us to analyze other T cell subsets for their usage of FcR γ . For this purpose, we examined the expression levels of TCR in T cells from $\zeta T/\zeta T$ mice and compared them with those in T cells from wild type mice. Since η is far less efficient in assembly and surface expression of TCR compared with ζ (20, 22), it is reasonable to assume that T cells whose TCR expression is resistant to the ζT mutation utilize FcR γ as a component of the TCR complex. Similar to thymocytes and peripheral T cells, TCR on CD3^{int}IL-2R β ^{hi} hepatic lymphocytes (30) were reduced in $\zeta T/\zeta T$ mice (Fig. 3 A). By contrast, surface TCR expression of r-IEL and DETC from $\zeta T/\zeta T$ mice remained higher than that of peripheral T cells, indicating the predominant usage of FcR γ in these cells (Fig. 3 B).

The present study demonstrates that in mice, certain subsets of T cells preferentially utilize FcR γ , of ζ family molecules, as a component of the TCR complex. These include CD8 $\alpha\alpha^+$ γ/δ^+ i-IEL, r-IEL, and DETC. It is of note that all these T cells express the TCR- γ/δ dimer and are local-

ized in epithelia. In addition, these cells express a higher level of TCR compared with peripheral T cells. These three subsets of γ/δ^+ T cells are thought to be phylogenically old and to play a part in the surveillance of body surfaces that are exposed to the environment (26, 31, 32). In this respect, it is of particular interest that these T cells commonly utilize FcR γ as a component of their TCR complexes. FcR γ has one signaling motif in the cytoplasmic portion, whereas ζ has three motifs, which are supposed to be generated by intramolecular duplication. Furthermore, FcR γ is also expressed in NK cells, which are proposed to be the ancestor of T cells (33). Taken together, FcR γ may be a prototype of ζ family molecules.

Differential usage of ζ and FcR γ by distinct T cell subsets likely represents differences in lineage and/or function of these cells. Recently, evidence supporting the idea that TCR with FcR γ is functionally distinct from TCR with ζ is increasing. First, in vitro kinase assay on immunoprecipitates of TCR complexes containing FcR γ displayed a distinct phosphorylation pattern from that of TCR complexes containing ζ in T hybridoma cells (17). Second, the cytoplasmic tail of ζ but not FcR γ can transduce signals mediated through Thy-1 molecules (18). Third, ζ is replaced by FcR γ in T cells from tumor-bearing mice as well as cancer patients, and the change is accompanied with immune incompetence including an impaired antitumor response (19). Concerning the functional difference between distinct T cell subsets, an interesting result was reported in mucosal immunity (34); γ/δ^+ i-IEL from mice orally immunized with antigen abrogates tolerance induced in recipient mice by oral administration of the same antigen when transferred, while α/β^+ i-IEL provides helper function for antibody production in vitro. These functional differences may reflect differential usage of ζ family molecules by γ/δ^+ and α/β^+ i-IEL. Further study should shed light on the physiological role of T cell subsets bearing TCR with different ζ family molecules.

We are grateful to Drs. R. Kubo, S. Tonegawa, C. Ra and M. Miyasaka for providing antibodies. We thank Dr. R. N. Germain, and Dr. Y. Ishida for critical reading of the manuscript, Ms. C. Ogata for technical assistance, and Ms. H. Ohno for secretarial assistance. ζT mice were established under collaboration with Dr. K. Rajewsky.

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture and the Life Science Research Project of Physical and Chemical Research (RIKEN).

Address correspondence to Takashi Saito, Division of Molecular Genetics, Center for Biomedical Science, Chiba University School of Medicine, 1-8-1 Inohana, Chuo-Ku, Chiba 260, Japan.

Received for publication 25 October 1993.

Note Added in Proof: After submitting this paper, Malissen et al. (36) reported a similar observation that TCR complexes of i-IEL contain FcR γ homodimers.

References

- Saito, T., A. Weiss, J. Miller, M.A. Norcross, and R.N. Germain. 1987. Specific antigen-Ia activation of transfected human T cells expressing murine T $\alpha\beta$ -human receptor complexes. *Nature (Lond.)* 325:125.
- Davis, M.M., and P.J. Bjorkman. 1988. T-cell antigen receptor genes and T-cell recognition. *Nature (Lond.)* 334:395.

3. Klausner, R.D., J. Lippincott-Schwartz, and J.S. Bonifacino. 1990. The T cell antigen receptor: insights into organelle biology. *Annu. Rev. Cell Biol.* 6:403.
4. Orloff, D.G., C. Ra, S.J. Frank, R.D. Klausner, and J.-P. Kinet. 1990. Family of disulfide-linked dimers containing the ζ and η chains of the T-cell receptor and the γ chain of Fc receptors. *Nature (Lond.)* 347:189.
5. Vivier, E., N. Rochet, J.P. Kochan, D.H. Presky, S.T. Schlossman, and P. Anderson. 1991. Structural similarity between Fc receptors and T cell receptors: expression of the γ -subunit of Fc ϵ RI in human T cells, natural killer cells and thymocytes. *J. Immunol.* 146:4263.
6. Koyasu, S., L. D'Adamio, A.R.N. Arulanandam, S. Abraham, L.K. Clayton, and E.L. Reinherz. 1992. T cell receptor complexes containing Fc ϵ RI γ homodimers in lieu of CD3 ζ and CD3 η components: a novel isoform expressed on large granular lymphocytes. *J. Exp. Med.* 175:203.
7. Minami, Y., A.M. Weissman, L.E. Samelson, and R.D. Klausner. 1987. Building a multichain receptor: synthesis, degradation, and assembly of the T-cell antigen receptor. *Proc. Natl. Acad. Sci. USA.* 84:2688.
8. Sussman, J.J., J.S. Bonifacino, J. Lippincott-Schwartz, A.M. Weissman, T. Saito, R.D. Klausner, and J.D. Ashwell. 1988. Failure to synthesize the T cell CD3- ζ chain: structure and function of a partial T cell receptor complex. *Cell.* 52:85.
9. Weissman, A.M., S.J. Frank, D.G. Orloff, M. Mercep, J.D. Ashwell, and R.D. Klausner. 1989. Role of the zeta chain in the expression of the T cell antigen receptor: genetic reconstitution studies. *EMBO (Eur. Mol. Biol. Organ.) J.* 8:3651.
10. Ashwell, J.D., and R.D. Klausner. 1990. Genetic and mutational analysis of the T-cell antigen receptor. *Annu. Rev. Immunol.* 8:139.
11. Irving, B.A., and A. Weiss. 1991. The cytoplasmic domain of the T cell receptor ζ chain is sufficient to couple to receptor-associated signal transduction pathways. *Cell.* 64:891.
12. Romeo, C., and B. Seed. 1991. Cellular immunity to HIV activated by CD4 fused to T cell or Fc receptor polypeptides. *Cell.* 64:1037.
13. Weiss, A. 1993. T cell antigen receptor signal transduction: a tale of tails and cytoplasmic protein-tyrosine kinases. *Cell.* 73:209.
14. Reth, M. 1989. Antigen receptor tail clue. *Nature (Lond.)* 338:383.
15. Wegener, A.-M., F. Letourneur, A. Hoeveler, T. Brocker, F. Luton, and B. Malissen. 1992. The T cell receptor/CD3 complex is composed of at least two autonomous transduction modules. *Cell.* 68:83.
16. Letourneur, F., and R.D. Klausner. 1992. Activation of T cells by a tyrosine kinase activation domain in the cytoplasmic tail of CD3 ϵ . *Science (Wash. DC)* 255:79.
17. Ohno, H., T. Aoe, C. Ra, T. Yamamoto, and T. Saito. 1993. TCR isoform containing the Fc receptor chain exhibit structural and functional differences from isoform containing CD3 ζ . *Int. Immunol.* 5:1403.
18. Letourneur, F., and R.D. Klausner. 1991. T-cell and basophil activation through the cytoplasmic tail of T-cell-receptor ζ family proteins. *Proc. Natl. Acad. Sci. USA.* 88:8905.
19. Mizoguchi, H., J.J. O'Shea, D.L. Longo, C.M. Loeffler, D.W. McVicar, and A.C. Ocha. 1992. Alterations in signal transduction molecules in T lymphocytes from tumor-bearing mice. *Science (Wash. DC)* 258:1795.
20. Ohno, H., T. Aoe, S. Taki, D. Kitamura, Y. Ishida, K. Rajewsky, and T. Saito. 1993. Developmental and functional impairment of T cells in mice lacking CD3 ζ chains. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4357.
21. Love, P.E., E.W. Shores, M.D. Johnson, M.L. Tremblay, E.J. Lee, A. Grinberg, S.P. Huang, A. Singer, and H. Westphal. 1993. T cell development in mice that lack the ζ chain of the T cell antigen receptor complex. *Science (Wash. DC)* 261:918.
22. Clayton, L.K., A. Diener, A. Lerner, A.G.D. Tse, S. Koyasu, and E.L. Reinherz. 1992. Differential regulation of T-cell receptor processing and surface expression affected by CD3 θ , an alternatively spliced product of the CD3 ζ/η gene locus. *J. Biol. Chem.* 267:26023.
23. Sudo, K., and K. Suzuki. 1987. Establishment of a new inbred nude strain, KSN. *Mouse News Lett.* 79:13.
24. Ishikawa, H., Y. Li, A. Abeliovich, S. Yamamoto, S.H.E. Kaufmann, and S. Tonegawa. 1993. Cytotoxic and interferon γ -producing activities of $\gamma\delta$ T cells in the mouse intestinal epithelium are strain dependent. *Proc. Natl. Acad. Sci. USA.* 90:8204.
25. Watanabe, H., K. Ohtsuka, M. Kimura, Y. Ikarashi, K. Ohmori, A. Kusumi, T. Ohteki, S. Seki, and T. Abo. 1992. Details of an isolation method for hepatic lymphocytes in mice. *J. Immunol. Methods.* 146:145.
26. Itohara, S., A.G. Farr, J.J. Lafaille, M. Bonneville, Y. Takagi, W. Haas, and S. Tonegawa. 1990. Homing of a $\gamma\delta$ thymocyte subset with homogeneous T-cell receptors to mucosal epithelia. *Nature (Lond.)* 343:754.
27. Ota, Y., T. Kobata, M. Seki, Y. Yagita, S. Shimada, Y.Y. Huang, Y. Takagi, Y., and K. Ohmura. 1992. Extrathymic origin of V γ 1/V δ 6 T cells in the skin. *Eur. J. Immunol.* 22:5958.
28. Miyake, K., I.L. Weissman, J.S. Greenberger, and P.W. Kincade. 1991. Evidence for a role of the integrin VLA-4 in lymphohemopoiesis. *J. Exp. Med.* 173:599.
29. Lefrancois, L. 1991. Extrathymic differentiation of intraepithelial lymphocytes: generation of a separate and unequal T-cell repertoire? *Immunol. Today.* 12:436.
30. Abo, T. 1993. Extrathymic pathways of T-cell differentiation: a primitive and fundamental immune system. *Microbiol. Immunol.* 37:247.
31. Janeway, C.A., Jr. 1988. Frontiers of the immune system. *Nature (Lond.)* 333:804.
32. Janeway, C.A., Jr., B. Jones, and A. Hayday. 1988. Specificity and function of T cells bearing $\gamma\delta$ receptors. *Immunol. Today.* 9:73.
33. Rodewald, H.-R., P. Moningeon, J.L. Lucich, C. Dosiou, P. Lopez, and E.L. Reinherz. 1992. A population of early fetal thymocytes expressing FcR2II/III contains precursors of T lymphocytes and natural killer cells. *Cell.* 69:139.
34. Fujihashi, K., T. Taguchi, W.K. Aicher, J.R. McGhee, J.A. Blustone, J.H. Eldridge, and H. Kiyono. 1992. Immunoregulatory functions for murine intraepithelial lymphocytes: γ/δ T cell receptor-positive (TCR $^+$) T cells abrogate oral tolerance, while α/β TCR $^+$ T cells provide B cell help. *J. Exp. Med.* 175:695.
35. Aoe, T., H. Ohno, S. Sonta, and T. Saito. 1993. Functional inactivation of the CD3 ζ chain in a T cell hybridoma by in vitro gene targeting. *Int. Immunol.* 5:725.
36. Malissen, M., A. Gillet, B. Rocha, J. Trucy, E. Vivier, C. Boyer, F. Kontgen, N. Brun, G. Mazza, E. Spanopoulou, et al. 1993. T cell development in mice lacking the CD3- ζ/η gene. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4347.