

Perini et al. Vol. 195, No. 7, April 1, 2002. Pages 907–918.

Due to a production error, the column heading of BETrka was missing. The corrected table appears below.

Table I. *Cell Death Induced by A β Peptides*

	BENTR-free		BEp75		BETrka	BEp75Trka
	24 h	48 h	24 h	48 h	24 h	24 h
Controls	5.9 \pm 2.6 ^a (18)	7.8 \pm 2.6 (18)	8.2 \pm 2.2 (62)	9.2 \pm 2.4 (20)	5.3 \pm 2.5 (3)	10.3 \pm 2.1 (3)
A β (25–35)	5.2 \pm 2.6 (12)	6.2 \pm 2.8 (12)	29.7 \pm 4.5 ^b (65)	34.0 \pm 4.8 ^b (14)	7.1 \pm 3.1 (5)	28.5 \pm 6.3 ^b (5)
A β (1–40)	5.6 \pm 2.3 (5)	5.5 \pm 2.8 (4)	27.6 \pm 2.3 ^b (5)	33.2 \pm 6.2 ^b (4)		
A β (1–42)	5.6 \pm 1.6 (5)	7.3 \pm 1.8 (4)	30.0 \pm 2.5 ^b (3)	30.9 \pm 2.2 ^b (4)		

Cells were treated with A β (25–35) (20 μ M), A β (1–40) (5.0 μ M), or A β (1–42) (5.0 μ M) and cell death was assessed by epifluorescence microscopy after 24 and 48 h.

^aThe values express percentages of cell death and are means \pm SD of the experiments indicated within brackets.

^b $P < 0.001$ with respect to the controls of the corresponding time point.